




## CLINICAL TREATMENT & MONITORING

### Diagnostic and Treatment Algorithm for Blastomycosis in Dogs or Cats



Antibody EIA test is not available for cats and Antibody ID test would be used instead.

\*Monitor urine antigen unless above the limit of quantification then monitor serum until urine antigen concentration is quantifiable.

^Mild-moderate disease could be defined as disease which can be treated on an out-patient basis and life-threatening disease requires hospitalization.

#### HEADQUARTERS


4705 Decatur Blvd. | Indianapolis, Indiana 46241, USA

888-841-8387



## CLINICAL TREATMENT & MONITORING

### Diagnostic and Treatment Algorithm for Histoplasmosis in Dogs or Cats




\*Monitor urine antigen unless above the limit of quantification then monitor serum until urine antigen concentration is quantifiable.

^Mild-moderate disease could be defined as disease which can be treated on an out-patient basis and life-threatening disease requires hospitalization.



## CLINICAL TREATMENT & MONITORING

### Diagnostic and Treatment Algorithm for Coccidioidomycosis (Valley Fever) in Dogs or Cats



Antibody EIA test is not available for cats and Antibody ID test would be used as the sole initial screening test.

#Use antigen test for treatment monitoring if positive at baseline. If not use antibody tests for monitoring.

^Mild-moderate disease could be defined as disease which can be treated on an out-patient basis and life-threatening disease requires hospitalization.

\*Fluconazole is the most commonly used oral treatment, but in vitro sensitivity data and clinical data from human medicine suggests that itraconazole should be considered as first-line treatment.

#### HEADQUARTERS


4705 Decatur Blvd. | Indianapolis, Indiana 46241, USA

888-841-8387



## CLINICAL TREATMENT & MONITORING

### Diagnostic and Treatment Algorithm for Cryptococcosis in Dogs or Cats



<sup>^</sup>Mild-moderate disease could be defined as disease which can be treated on an out-patient basis and life-threatening disease requires hospitalization.

#### HEADQUARTERS


4705 Decatur Blvd. | Indianapolis, Indiana 46241, USA

888-841-8387



**CLINICAL TREATMENT & MONITORING**

**Diagnostic and Treatment Algorithm for Systemic Aspergillosis in Dogs or Cats**




\*Use whichever sample had the highest initial antigen concentration for treatment monitoring.

^Mild-moderate disease could be defined as disease which can be treated on an out-patient basis and life-threatening disease requires hospitalization.



## CLINICAL TREATMENT & MONITORING

### Diagnostic and Treatment Algorithm for Sinonasal Aspergillosis in Dogs or Cats



#### HEADQUARTERS

4705 Decatur Blvd. | Indianapolis, Indiana 46241, USA

Version 1 | May 2021

888-841-8387

[miravistavets.com](http://miravistavets.com)